

$p = 0.01$). In a subgroup analysis, patients who underwent transfemoral TAVR had the lowest in-hospital mortality, post-procedure length of stay, need for blood transfusion, and cost of hospitalization, compared with transapical TAVR and SAVR.

Over the 11-month period in the NRD, a total of 446 patients with ESRD had undergone an aortic valve replacement; 118 had TAVR (73 transfemoral and 26 transapical), and 347 had SAVR. Of these, 104 (23.32%) patients had a readmission within a 30-day period after discharge. No differences between the 30-day readmission rates were observed after TAVR and SAVR. The median times to readmission, length of stay, and in-hospital mortality during the 30-day readmission were also similar for the patient cohorts.

Our findings suggested that the patients with ESRD are at higher risk of mortality (13.21%) and complications after aortic valve replacement procedures. Almost 1 of 4 patients were readmitted within 30 days. Despite selecting older ESRD patients, TAVR resulted in similar mortality and 30-day readmission rates when compared to SAVR, but had a shorter length of stay and lower hospitalization costs. The likelihood of discharge to home was higher after TAVR. Transfemoral TAVR appeared to be the safest and the most effective treatment for these patients, sustaining the least mortality.

Viraj Bhise, MBBS, MPH
Pushkar Kanade, MD, MPH
Ghanshyam P.S. Shantha, MD, MPH
Prakash Balan, MD
Tom C. Nguyen, MD
Pranav Loyalka, MD
Biswajit Kar, MD
Anthony Estrera, MD

Richard W. Smalling, MD, PhD

*Abhijeet Dhoble, MD, MPH

*Department of Cardiology
University of Texas Health Science Center
6431 Fannin Street, MSB 1.224
Houston, Texas, 77030

E-mail: abhijeet.dhoble@uth.tmc.edu

<http://dx.doi.org/10.1016/j.jacc.2017.01.049>

Please note: Dr. Nguyen is a consultant for Edwards Lifesciences. All other authors have reported that they have no relationships relevant to the contents of this paper to disclose.

REFERENCES

- Ohno Y, Attizzani GF, Barbanti M, et al. Transcatheter aortic valve replacement for severe aortic stenosis patients undergoing chronic dialysis. *J Am Coll Cardiol* 2015;66:93-4.
- Houchens RL, Ross DN, Elixhauser A, Jiang J. Nationwide inpatient sample redesign: final report. Rockville, MD: Agency for Healthcare Research and Quality, 2014.

3. Healthcare Cost and Utilization Project. NRD description of data elements. Rockville, MD: Agency for Healthcare Research and Quality, 2015.

4. Leuven E, Sianesi B. PSMATCH2: Stata module to perform full Mahalanobis and propensity score matching, common support graphing, and covariate imbalance testing. Boston, MA: Statistical Software Components, Boston College, 2015.

Combination Therapy With Ceftriaxone and Lansoprazole, Acquired Long QT Syndrome, and Torsades de Pointes Risk



We read with great interest the paper by Lorberbaum et al. (1) recently published in the *Journal*, in which evidence was provided that ceftriaxone plus lansoprazole combination therapy (CFX+LSZ) was unexpectedly associated with an increased risk of acquired long QT syndrome (aLQTS); thus, it is also theoretically associated with Torsades de Pointes (TdP) and sudden cardiac death. We report on 2 clinical observations that possibly further support these findings.

First, we recently observed a patient affected with pneumonia who specifically developed aLQTS after CFX+LSZ administration.

On admission, the patient's electrocardiogram (ECG) showed a normal corrected QT interval (QTc; 422 ms). Ceftriaxone administration was started (2 g/day), and a second ECG performed 3 days later did not reveal any significant change in the QTc (417 ms). On day 8, the patient reported epigastralgia, and lansoprazole (30 mg/day) was added to therapy. After 2 days of the combined treatment, QTc prolongation developed (475 ms), despite no changes in electrolytes, echocardiography findings, or further drug use. Lansoprazole was stopped and replaced with pantoprazole, whereas ceftriaxone was continued. Two days later, QTc normalized with restoration of baseline values (424 ms).

Second, in a cohort of unselected TdP patients, we observed that CFX+LSZ was assumed by a non-negligible percentage of subjects.

Since 2008, we have prospectively recruited patients who experienced TdP, independent of ongoing therapies and concomitant diseases. To date, 40 consecutive patients have been recruited. As expected (2), a high prevalence of recognized QT-prolonging factors of acquired origin were present (on average >4), including mainly cardiac diseases (82%), electrolyte imbalances (70%),

QT-prolonging medications (57%, globally considered), and anti-Ro/Sjogren's syndrome-related antigen A antibodies (52%) (3).

In this cohort, we found that 2 patients (5%) were receiving CFX+LSZ when TdP developed. Notably, this percentage was identical to that observed with levofloxacin, clarithromycin, or promazine (5% each) and was higher than that seen with ciprofloxacin, sotalolol, or haloperidol (2%), which all represent well-recognized QT-prolonging medications (4). Only amiodarone (27%) and citalopram, fluconazole, or sertraline (7%) were more frequently administered.

Although follow-up studies are required to confirm the findings of Lorberbaum et al. (1), our observations seem to be in agreement with the view that CFX+LSZ may represent a currently overlooked contributing factor that increases the TdP risk, via specific (not class-related) QT-prolonging effects exerted by these 2 molecules when combined.

The potential harmfulness of this association should be carefully kept in mind, particularly in the presence of other recognized QT-prolonging factors. Both drugs are used worldwide, but they are presently considered as having neutral effects on ventricular repolarization.

***Pietro Enea Lazznerini, MD**
Iacopo Bertolozzi, MD
Marco Rossi, PharmD
Pier Leopoldo Capecchi, MD
Franco Laghi-Pasini, MD

*Department of Medical Sciences, Surgery
and Neurosciences
University of Siena
Viale Bracci 14
53100 Siena
Italy

E-mail: lazznerini7@unisi.it

<http://dx.doi.org/10.1016/j.jacc.2016.11.090>

Please note: The authors have reported that they have no relationships relevant to the contents of this paper to disclose.

REFERENCES

1. Lorberbaum T, Sampson KJ, Chang JB, et al. Coupling data mining and laboratory experiments to discover drug interactions causing QT prolongation. *J Am Coll Cardiol* 2016;68:1756-64.
2. Drew BJ, Ackerman MJ, Funk M, et al. Prevention of torsade de pointes in hospital settings: a scientific statement from the American Heart Association and the American College of Cardiology Foundation. *J Am Coll Cardiol* 2010;55:934-47.
3. Lazznerini PE, Yue Y, Srivastava U, et al. Arrhythmogenicity of anti-Ro/SSA antibodies in patients with torsades de pointes. *Circ Arrhythm Electrophysiol* 2016;9:e003419.
4. Woosley RL, Romero KA. Crediblemeds.org QT drugs List, 2016. AZCERT, Inc. Available at: <https://crediblemeds.org>. Accessed October 18, 2016.

REPLY: Combination Therapy With Ceftriaxone and Lansoprazole, Acquired Long QT Syndrome and Torsades de Pointes Risk



We thank Dr. Lazznerini and colleagues for sharing their clinical observations supporting the drug-drug interaction (DDI) findings in our recent paper (1). In isolation, their observations could be dismissed as idiosyncratic by skeptics, but data science—the application of rigorous analytical methods to large datasets—provides the context needed to explain these effects across large patient populations. The integration of adverse event report mining, electronic health record corroboration, electrophysiology experiments, and now these case studies paints a clear picture of a DDI that could not have been identified using traditional surveillance approaches (2).

We have further capitalized on the promise of health data science by participating in Observational Health Data Sciences and Informatics (OHDSI), an international network of >140 researchers in >20 countries performing large-scale analyses on distributed databases that in total comprise >600 million patients (3). Each participating site converts their patient data to a common data model that enables analyses created at one location to be replicated across the entire network while maintaining patient privacy. This open and international collaboration has developed open source software tools for data exploration and evidence generation, conducted multivariate analyses to predict drug side effects, and improved definition of the heterogeneity in treatment pathways (4,5). The latter study was performed on an underlying population of >250 million patients from 11 data sources in 4 countries and took 20 days from conception to results from 7 sources, highlighting the efficiency with which these analyses can be performed at scale.

In the case of ceftriaxone and lansoprazole, we used a combination of public and private clinical data to identify the interaction. Through collaborations like OHDSI, these data are available to all researchers with interests in large-scale retrospective studies. We hope that the current study motivates others to participate in these new data science initiatives.

Tal Lorberbaum, MA

***Nicholas P. Tatonetti, PhD**

*Department of Biomedical Informatics
Columbia University
622 West 168th Street, PH20
New York, New York 10032

E-mail: nick.tatonetti@columbia.edu

<http://dx.doi.org/10.1016/j.jacc.2016.12.042>

Please note: Dr. Tatonetti is a compensated advisor to Advera Health, Inc. Mr. Lorberbaum has reported that he has no relationships relevant to the contents of this paper to disclose.

REFERENCES

1. Lorberbaum T, Sampson KJ, Chang JB, et al. Coupling data mining and laboratory experiments to discover drug interactions causing QT prolongation. *J Am Coll Cardiol* 2016;68:1756-64.
2. Lorberbaum T, Sampson KJ, Woosley RL, et al. An integrative data science pipeline to identify novel drug interactions that prolong the QT interval. *Drug Saf* 2016;39:433-41.
3. Observational Health Data Sciences and Informatics. Available at: <https://www.ohdsi.org/>. Accessed March 3, 2017.
4. Ryan PB, Madigan D, Stang PE, et al. Medication-wide association studies. *CPT Pharmacometrics Syst Pharmacol* 2013;2:e76.
5. Hripscak G, Ryan PB, Duke JD, et al. Characterizing treatment pathways at scale using the OHDSI network. *Proc Natl Acad Sci USA* 2016;113:7329-36.

Vitamin E May Protect Against Contrast-Induced Acute Kidney Injury



McCullough et al. (1) reviewed the pathophysiology and treatment options for contrast-induced acute kidney injury (CI-AKI). They stated that no effective adjunctive pharmaceutical had been demonstrated that either prevented or treated CI-AKI. However, they also suggested that of the agents being investigated, statins were the most promising. We would like to point out that strong evidence has also emerged regarding the effect of vitamin E against CI-AKI, which was not mentioned in their review.

Three randomized placebo-controlled trials found that vitamin E significantly prevented CI-AKI, with point estimates ranging from 52% to 75% for the decrease in the incidence of CI-AKI (2-4). All participants had chronic kidney disease and had been subjected to coronary catheterization or angiography. The latest trial reported CI-AKI cases in 14.1% of the placebo group, but in only 6.7% of the vitamin E group, which corresponded to 7.4% of participants benefitting from the vitamin, with a number needed to treat (NNT) of 13.5 (4). The 2 earlier studies found NNTs of 5.8 (2) and 10.6 (3). In each study, approximately one-half of the patients were on statin therapy; therefore, the effects of vitamin E might have also been beneficial in addition to statins.

Two of the vitamin E trials were carried out in Thailand (2,3), and 1 was carried out in Iran (4). Thus,

it is not known whether the findings can be directly generalized to Western countries. Even if the positive findings might only be applicable to less developed countries, the findings are important for the populations of such countries.

Vitamin E is an essential nutrient, and therefore, its potential benefit in preventing CI-AKI is interesting. Furthermore, vitamins E and C may interact. Vitamin E decreased total mortality in male smokers aged older than 65 years if their dietary vitamin C intake level was high, but not if their vitamin C intake was low (5). Thus, a large factorial trial seems warranted to examine the effect of statins and vitamins E and C, and their combinations to discover the optimal protocol to prevent CI-AKI.

*Harri Hemilä, MD, PhD

Yousef Rezaei, MD

*Department of Public Health

University of Helsinki

POB 20, Tukholmankatu 8B

Helsinki FIN-00014

Finland

E-mail: harri.hemila@helsinki.fi

<http://dx.doi.org/10.1016/j.jacc.2016.11.091>

Please note: Both authors have reported that they have no relationships relevant to the contents of this paper to disclose.

REFERENCES

1. McCullough PA, Choi JP, Feghali GA, et al. Contrast-induced acute kidney injury. *J Am Coll Cardiol* 2016;68:1465-73.
2. Tasanarong A, Piyayotai D, Thitiarchakul S. Protection of radiocontrast induced nephropathy by vitamin E (alpha tocopherol): a randomized controlled pilot study. *J Med Assoc Thai* 2009;92:1273-81.
3. Tasanarong A, Vohakiat A, Hutayanon P, et al. New strategy of alpha- and gamma-tocopherol to prevent contrast-induced acute kidney injury in chronic kidney disease patients undergoing elective coronary procedures. *Nephrol Dial Transplant* 2013;28:337-44.
4. Rezaei Y, Khademvatani K, Rahimi B, et al. Short-term high-dose vitamin E to prevent contrast medium-induced acute kidney injury in patients with chronic kidney disease undergoing elective coronary angiography: a randomized placebo-controlled trial. *J Am Heart Assoc* 2016;5:e002919.
5. Hemilä H, Kaprio J. Modification of the effect of vitamin E supplementation on the mortality of male smokers by age and dietary vitamin C. *Am J Epidemiol* 2009;169:946-53.

REPLY: Vitamin E May Protect Against Contrast-Induced Acute Kidney Injury



We appreciate the comments from Drs. Hemilä and Rezaei concerning small randomized trials of short-term vitamin E for the prevention of contrast-induced acute kidney injury (CI-AKI). In the trial